

## 参考文献

- 1 郭素淑, 李曙光. 淡色花岗岩的岩石学和地球化学特征及其成因. *地学前缘*, 2007, 14(6): 290—298
- 2 Le Fort P, Cuney C, Deniel C, et al. Crustal generation of the Himalayan leucogranites. *Tectonophysics*, 1987, 134(1-3):39—57[[doi](#)]
- 3 Debon F, Le Fort P, Sheppard S M F, et al. The four pluton belts of the Transhimalaya-Himalaya: A chemical, mineralogical, isotopic, and chronological synthesis along a Tibet-Nepal section. *J Petrol*, 1986, 27(1): 219—250
- 4 Guillot S, Le Fort P. Geochemical constraints on the bimodal origin of High Himalayan leucogranites. *Lithos*, 1995, 35(3-4): 221—234[[doi](#)]
- 5 Harrison T M, Ryerson F J, Le Fort P, et al. A Late Miocene-Pliocene origin for the central Himalayan inverted metamorphism. *Earth Planet Sci Lett*, 1997, 146(1-2): E1—E7[[doi](#)]
- 6 Barbey P, Brouand M, Le Fort P, et al. Granite-migmatite genetic link: The example of the Manaslu granite and Tibetan slab migmatites in central Nepal. *Lithos*, 1996, 38(1-2): 63—79[[doi](#)]
- 7 Visonà D, Lombardo B. Two-mica and tourmaline leucogranites from the Everest-Makalu region (Nepal-Tibet). Himalayan leucogranite genesis by isobaric heating? *Lithos*, 2002, 62(3-4): 125—250[[doi](#)]
- 8 Inger S, Harris N B W. Tectonothermal evolution of the High Himalayan Crystalline Sequence, Langtang Valley, northern Nepal. *J Metamorph Geol*, 1992, 10(3): 439—452[[doi](#)]
- 9 Inger S, Harris N. Geochemical constraints on leucogranite magmatism in the Langtang Valley, Nepal Himalaya. *J Petrol*, 1993, 34(2): 345—368
- 10 Harris N, Ayres M, Massey J. Geochemistry of granitic melts produced during the incongruent melting of muscovite: Implications for the extraction of Himalayan leucogranite magmas. *Geophys Res*, 1995, 100(B8): 15767—15777[[doi](#)]
- 11 Searle M P, Parrish R R, Hodges K V, et al. Shisha Pangma leucogranite, south Tibetan Himalaya: Field relations, geochemistry, age, origin, and emplacement. *J Geol*, 1997, 105(3): 295—317
- 12 张金阳, 廖群安, 李德威. 西藏定结地区高喜马拉雅淡色花岗岩的地球化学特征与岩浆源区研究. *地质科技情报*, 2003, 22(3): 9—14
- 13 Ayres M, Harris N. Ree fractionation and Nd-isotope disequilibrium during crustal anatexis: Constraints from Himalayan leucogranites. *Chem Geol*, 1997, 139(1-4): 249—269[[doi](#)]
- 14 Scaillet B, France-Lanord C, Le Fort P. Badrinath-Gangotri plutons (Garhwal, India): Petrological and geochemical evidence for fractionation processes in a high Himalayan leucogranite. *J Volcan Geothermal Res*, 1990, 44(1-2): 163—188[[doi](#)]
- 15 杨晓松, 金振民. 西藏亚东淡色花岗岩 Rb-Sr 和 Sm-Nd 同位素研究——关于其年龄和源岩的证据. *地质论评*, 2001, 47(3): 294—300
- 16 张金阳, 廖群安, 李德威, 等. 藏南萨迦拉纳岗日淡色花岗岩特征及与变质核杂岩的关系. *地球科学——中国地质大学学报*, 2003, 28(6): 695—701
- 17 张宏飞, Harris N, Parrish R, 等. 北喜马拉雅淡色花岗岩地球化学区域对比、岩石成因及其构造意义. *地球科学——中国地质大学学报*, 2005, 30(3): 275—288
- 18 陈斌, 翟明国, 邵济安. 太行山北段中生代岩基的成因和意义: 主要和微量元素地球化学证据. *中国科学D辑: 地球科学*, 2002, 32(11): 898—907
- 19 Nabelek P I, Russ-Nabelek C, Denison J R. The generation and crystallization conditions of the Proterozoic Harney Peak leucogranite, Black Hills, South Dakota, USA: petrologic and geochemical constraints. *Contrib Mineral Petrol*, 1992, 110(2-3): 173—191[[doi](#)]
- 20 Nabelek P I, Bartlett C D. Petrologic and geochemical links between the post-collisional Proterozoic Harney Peak leucogranite, South Dakota, USA, and its source rocks. *Lithos*, 1998, 45(1-4): 71—85[[doi](#)]
- 21 van de Fliert D T, Jung S, Masberg P, et al. Lower crustal melting and the role of open-system processes in the genesis of syn-orogenic quartz diorite-granite-leucogranite associations: Constraints from Sr-Nd-O isotopes from the Bandombaai complex, Namibia. *Lithos*, 2003, 67(3-4): 205—226[[doi](#)]
- 22 Pressley R A, Brown M. The Phillips Pluton, Maine, USA: Evidence of heterogeneous crustal sources and implications for granite ascent and emplacement mechanisms in convergent orogens. *Lithos*, 1999, 46(3): 335—366[[doi](#)]
- 23 Solar G S, Brown M. Petrogenesis of migmatites in Maine, USA: Possible source of prealuminous leucogranite in plutons? *J Petrol*, 2001, 42(4): 789—823[[doi](#)]
- 24 Braun I, Raith M, Ravinera Kumar G R. Dehydration-melting phenomena in Leptynitic gneisses and the generation of leucogranites: A case study from the Kerala Khondalite belt, Southern India. *J Petrol*, 1996, 37(6): 1285—1305[[doi](#)]
- 25 郭素淑, 李曙光. 华北克拉通东南缘古元古代变质和岩浆事件的锆石 SHRIMP U-Pb 年龄. *中国科学D辑: 地球科学*, 2009, 39(5)
- 26 Liu Y C, Wang A D, Rolfo F, et al. Geochronological and petrological constraints on Palaeoproterozoic granulite facies metamorphism in southeastern margin of the North China Craton. *J Metamorph Geol*, 2009, 27(2): 125—138[[doi](#)]
- 27 许文良, 王清海, 杨德彬, 等. 蚌埠荆山“混合花岗岩”SHRIMP 锆石 U-Pb 定年及其地质意义. *中国科学D辑: 地球科学*, 2004, 34(5): 423—428
- 28 尚瑞钧, 严阵, 等. 秦巴花岗岩. 武汉: 中国地质大学出版社, 1998
- 29 钱春超, 路玉林, 刘丽利. 大别山超高压变质带燕山期花岗岩地球化学特征及成因探讨. *中国地质*, 2004, 31(2): 147—154

- 30 Zhang H F, Gao S, Zhong Z Q, et al. Geochemical and Sr-Nd-Pb isotopic compositions of Cretaceous granitoids: Constraints on tectonic framework and crustal structure of the Dabieshan ultrahigh-pressure metamorphic belt, China. *Chem Geol*, 2002, 186(3-4): 281—299[[doi](#)]
- 31 Xu H J, Ma C Q, Ye K. Early cretaceous granitoids and their implications for the collapse of the Dabie orogen, eastern China: SHRIMP zircon U-Pb dating and geochemistry. *Chem Geol*, 2007, 240(3-4): 238—259
- 32 徐小军, 赵子福, 郑永飞, 等. 大别造山带天柱山燕山期中酸性岩浆岩元素和同位素地球化学研究. *岩石学报*, 2005, 21(3): 607—622
- 33 马昌前, 杨坤光, 许长海, 等. 大别山中生代钾质岩浆作用与超高压变质地体的剥露机理. *岩石学报*, 1999, 15(3): 379—395
- 34 马昌前, 杨坤光, 明厚利, 等. 大别山中生代地壳从挤压转向伸展的时间: 花岗岩的证据. *中国科学 D 辑: 地球科学*, 2003, 33(9): 817—827
- 35 Chen B, Jahn Bor-ming, Wei C J. Petrogenesis of Mesozoic granitoids in the Dabie UHP complex. *Lithos*, 2002, 60(1-2): 67—88[[doi](#)]
- 36 Xie Z, Zheng Y F, Zhao Z F, et al. Mineral isotope evidence for the contemporaneous process of Mesozoic granite emplacement and gneiss metamorphism in the Dabie orogen. *Chem Geol*, 2006, 231(3): 214—235[[doi](#)]
- 37 齐瑞荣, 黄增保, 金霞. 甘肃北山大石山 A 型花岗岩体的地球化学特征及构造意义. *岩石矿物学杂志*, 2006, 25(2): 90—96
- 38 许保良, 阎国翰, 徐振邦, 等. 冀北燕山期三个系列花岗质岩石的地球化学特征及其成因学意义. *岩石学报*, 1999, 15(2): 208—216
- 39 朱金初, 张佩华, 谢才富, 等. 南岭西段花山-姑婆山 A 型花岗质杂岩带: 岩石学、地球化学和岩石成因. *地质学报*, 2006, 80(4): 529—542
- 40 顾晟彦, 华仁民, 戚华文. 广西花山-姑婆山燕山期花岗岩的地球化学及成因研究. *岩石矿物学杂志*, 2006, 25(2): 97—109
- 41 邱检生, 胡建, 王孝磊, 等. 广东河源白石岗岩体: 一个高分异的 I 型花岗岩. *地质学报*, 2005, 79(4): 503—514
- 42 Montel J M. A model for monazite/melt equilibrium and application to the generation of granitic magmas. *Chem Geol*, 1993, 110(1-3): 127—146[[doi](#)]
- 43 Li S G, Xiao Y L, Liu D L, et al. Collision of the North China and Yangtz Blocks and formation of coesite-bearing eclogites: Timing and processes. *Chem Geol*, 1993, 109(1-4): 89—111[[doi](#)]
- 44 郑永飞. 超高压变质与大陆碰撞研究进展: 以大别-苏鲁造山带为例. *科学通报*, 2008, 53(18): 2129—2152
- 45 Zhao Z F, Zheng Y F, Wei C S, et al. Post-collisional granitoids from the Dabie orogen in China: Zircon U-Pb age, element and O isotope evidence for recycling of subducted continental crust. *Lithos*, 2007, 93(3-4): 248—272 [doi]
- 46 Chemenda A I, Burg J P, Mattauer M. Evolutionary model of the Himalaya-Tibet system: Geopoeem based on new modeling, geological and geophysical data. *Earth Planet Sci Lett*, 2000, 174: 397—409 [doi]
- 47 Westaway R. Crustal volume balance during the India-Eurasia collision and altitude of the Tibetan plateau: A working hypothesis. *J Geophys Res*, 1995, 100(B8): 15173—15192[[doi](#)]
- 48 Zhao W L, Morgan W J. Uplift of the Tibetan Plateau. *Tectonics*, 1985, 4(4): 359—369[[doi](#)]
- 49 Chung S L, Chu M F, Zhang Y Q. Tibetan tectonic evolution inferred from spatial and temporal variations in post-collisional magmatism. *Earth Sci Rev*, 2005, 68(3-4): 173—196 [doi]
- 50 Rapp R P, Shimizu N, Norman M D, et al. Reaction between slab-derived melts and peridotite in the mantle wedge: Experimental constraints at 3.8 GPa. *Chem Geol*, 1999, 160(4): 335—356[[doi](#)]
- 51 Harrison T M, Lovera O M, Grove M. New insights into the origin of two contrasting Himalayan granite belts. *Geology*, 1997, 25(10): 899—902[[doi](#)]